

Patient information from BMJ

Last published: May 02, 2025

Whiplash

Whiplash is a neck injury that happens when the head is suddenly and forcefully jolted. It usually gets better on its own within a few weeks or months. But simple treatments may help with the pain and speed up your recovery.

What is whiplash?

Whiplash is a neck injury that can happen if your head is suddenly jolted backward and forward in a whip-like movement. When this happens the muscles, ligaments, and tendons in your neck stretch more than normal and may be sprained. Ligaments and tendons are tissues that connect your bones and muscles together.

The most common cause of whiplash is being in a vehicle accident. Other causes of whiplash include:

- falls (especially in older people)
- getting hurt in a physical fight
- sports injuries.

What are the symptoms of whiplash?

The main symptoms of whiplash are neck pain and neck stiffness. But often people don't feel these symptoms at the time of their injury. The pain and stiffness will probably be worse a day or two afterward.

Other symptoms of whiplash you may experience include:

- Headaches
- Pain in your upper back, arms, or shoulder
- Pain in your lower back
- Trouble concentrating
- Dizziness
- Ringing in your ears (tinnitus)

Whiplash

- Blurred vision.

Your doctor will ask you questions about your injury and how it happened. This will help them better understand if your symptoms are because of whiplash. They'll also want to do a physical exam.

A severe whiplash injury can cause more serious problems that need hospital treatment. You should tell your doctor if you notice:

- any weakness in your limbs
- numbness, tingling, or pins and needles in your limbs
- you're losing control of your bladder or bowels.

These could be signs of an injury to the nerves in your spine.

Most people with whiplash won't need any scans. But your doctor might recommend a CT scan of your spine if they think you have a more serious injury.

What are the treatment options for whiplash?

Most cases of whiplash can be treated at home with medications to target pain and stiffness. But if your doctor suspects you've injured your spine, then you will need to stay in the hospital for treatment.

You might feel as though you should rest and move as little as possible after whiplash. But it's important that you carry on with your normal activities as much as possible for your recovery.

You should also avoid wearing a neck brace. This stops your neck moving freely, and won't help your pain and stiffness.

Medications

Pain medications can help with the pain and stiffness from whiplash. Your doctor will usually recommend you try ibuprofen or naproxen first. These are medications known as **NSAIDs** (nonsteroid anti-inflammatory drugs).

NSAIDs work to target both pain and inflammation (swelling). But they can cause side effects in some people, like irritation of your stomach lining. So it's important that you follow your doctor or pharmacist's advice about taking this medication.

If NSAIDs don't help enough, your doctor might suggest a course of stronger pain relievers called **opioids**. These medications are only recommended for short periods. This is because there is a chance people can become dependent on them if they're taken for longer.

Physical therapy

Your doctor will recommend that you keep active as much as possible after your injury. But if your symptoms don't get better after 4-6 weeks, they may suggest you try physical therapy. This could be helpful if you're dealing with pain because you've been in a vehicle accident.

What happens next?

Whiplash usually gets better within a few weeks or months. In very mild cases, you may even start feeling better after a few days.

You can usually manage whiplash at home with pain relievers and follow-up with your doctor every few weeks. But you will need to stay in the hospital if you have a more serious injury.

It's important to keep up with your normal activities as much as possible, even if you're feeling pain and discomfort. This will help speed up your recovery.

If your symptoms don't seem to be getting better, or they're getting worse, then speak with your doctor. In some people, the pain and stiffness from whiplash can be chronic (i.e., long-lasting). Chronic pain lasts for more than 3 months. Your doctor will be able to give you more advice about how to manage this type of pain.

If you notice any **new** neurologic signs or symptoms, seek **immediate** medical attention. These can include:

- weakness in your limbs
- numbness, tingling, or pins and needles in your limbs
- losing control of your bladder or bowels.

The patient information from *BMJ Best Practice* is regularly updated. The most recent version of Best Practice can be found at bestpractice.bmj.com. This information is intended for use by health professionals. It is not a substitute for medical advice. It is strongly recommended that you independently verify any interpretation of this material and, if you have a medical problem, see your doctor.

Please see BMJ's full terms of use at: bmj.com/company/legal-information. BMJ does not make any representations, conditions, warranties or guarantees, whether express or implied, that this material is accurate, complete, up-to-date or fit for any particular purposes.

© BMJ Publishing Group Ltd 2025. All rights reserved.

What did you think about this patient information guide?

Complete the [online survey](#) or scan the QR code to help us to ensure our content is of the highest quality and relevant for patients. The survey is anonymous and will take around 5 minutes to complete.

